



PTFE Chemical Resistance

This chart is intended to be used as a general guide only. Since each rating is for ideal conditions, all factors affecting chemical resistance must be considered.

| Acetaldehyde | Е | Bromine | * | Cresol | Е |
|-------------------------|---|----------------------------|---|---------------------------|---|
| Acetamide, Sat. | Е | Bromobenzene | Е | Cyclohexane | Е |
| Acetic Acid, 5% | Е | Bromoform | Е | Decalin | Е |
| Acetic Acid, 50% | Е | Butadiene | Е | o-Dichlorobenzene | Е |
| Acetone | Е | n-Butyl Acetate | Е | p-Dichlorobenzene | Е |
| Acetonitrile | E | n-Butyl Alcohol | Е | Diethyl Benzene | Е |
| Acrylonitrile | Е | sec-Butyl Alcohol | Е | Diethyl Ether | Е |
| Adipic Acid | Е | tert-Butyl Alcohol | Е | Diethyl Ketone | Е |
| Alanine | Е | Butyric Acid | Е | Diethyl Malonate | Е |
| Allyl Alcohol | Е | Calcium Hydroxide, Conc. | Е | Diethylene Glycol | Е |
| Aluminum Hydroxide | Е | Calcium Hypochlorite, Sat. | Е | Dimethyl Formamide | Е |
| Aluminum Salts | Е | Carbazole | Е | Dimethylsulfoxide | Е |
| Amino Acids | Е | Calcium Hydroxide, Conc. | Е | 1,4-Dioxane | Е |
| Ammonia | Е | Calcium Hypochlorite, Sat. | Е | Dipropylene Glycol | Е |
| Ammonium Acetate, Sat. | Е | Carbazole | Е | Ether | Е |
| Ammonium Glycolate | Е | Carbon Disulfide | Е | Ethyl Acetate | Е |
| Ammonium Hydroxide 5% | Е | Carbon Tetrachloride | Е | Ethyl Alcohol (absolute) | Е |
| Ammonium, Hydroxide 30% | Е | Cedarwood Oil | Е | Ethyl Alcohol, 40% | Е |
| Ammonium Oxalate | Е | Cellosolve Acetate | Е | Ethyl Benzene | Е |
| Ammonium Salts | Е | Chlorine, 10% in Air | Е | Ethyl Benzoate | Е |
| n-Amyl Acetate | Е | Chlorine, 10% (Moist) | Е | Ethyl Butyrate | Е |
| Amyl Chloride | Е | Chloroacetic Acid | Е | Ethyl Chloride | Е |
| Aniline | Е | p-Chloroacetophenone | Е | Ethyl Cyanoacetate | Е |
| Benzaldehyde | Е | Chloroform | Е | Ethyl Lactate | Е |
| Benzene | Е | Chromic Acid, 10% | Е | Ethylene Chloride, Liquid | Е |
| Benzoic Acid, Sat. | Е | Chromic Acid, 50% | Е | Ethylene Glycol | Е |
| Benzyl Acetate | Е | Cinnamon Oil | Е | Ethylene Oxide | Е |
| Benzyl Alcohol | Е | Citric Acid, 10% | Е | Fluorides | Е |

^{* =} Moderate Attack or appreciable absorption. Material will have limited life.

E = Excellent





PTFE Chemical Resistance

This chart is intended to be used as a general guide only. Since each rating is for ideal conditions, all factors affecting chemical resistance must be considered.

| Fluorine | * | Methyl Ethyl Ketone | Е | Silver Acetate | Е |
|-------------------------|---|----------------------------|---|-------------------------------|---|
| Formaldehyde, 10% | Е | Methyl Isobutyl Ketone | Е | Silver Nitrate | Е |
| Formaldehyde, 40% | Е | Methyl Propyl Ketone | Е | Sodium Acetate, Sat. | Е |
| Formic Acid, 3% | Е | Methylene Chloride | Е | Sodium Hydroxide, 1% | Е |
| Formic Acid, 50% | Е | Mineral Oil | Е | Sodium Hydroxide 50%-Sat. | Е |
| Formic Acid, 98-100% | Е | Nitric Acid, 1-10% | Е | Sodium Hypochlorite, 15% | Е |
| Fuel Oil | Е | Nitric Acid, 50% | Е | Stearic Acid, Crystals | Е |
| Gasoline | Е | Nitric Acid, 70% | Е | Sulfuric Acid, 1-6% | Е |
| Glacial Acetic Acid | Е | Nitrobenzene | Е | Sulfuric Acid, 20% | Е |
| Glycerin | Е | n-Octane | Е | Sulfuric Acid, 60% | Е |
| n-Heptane | Е | Orange Oil | Е | Sulfuric Acid, 98% | Е |
| Hexane | Е | Ozone | Е | Sulfuric Dioxide, Liq., 46psi | Е |
| Hydrochloric Acid, 1-5% | Е | Perchloric Acid | * | Sulfuric Dioxide, wet or dry | Е |
| Hydrochloric Acid, 20% | Е | Perchloroethylene | Е | Sulfur Salts | Е |
| Hydrochloric Acid, 35% | Е | Phenol, Crystals | Е | Tartaric Acid | Е |
| Hydrofluoric Acid, 4% | Е | Phosphoric Acid, 1-5% | Е | Tetrahydrofuran | Е |
| Hydrofluoric Acid, 48% | Е | Phosphoric Acid, 85% | Е | Thionyl Chloride | Е |
| Hydrogen Peroxide, 3% | Е | Pine Oil | Е | Toluene | Е |
| Hydrogen Peroxide, 30% | Е | Potassium Hydroxide, 1% | Е | Tributyl Citrate | Е |
| Hydrogen Peroxide, 90% | Е | Potassium Hydroxide, Conc. | Е | Trichloroethane | Е |
| Isobutyl Alcohol | Е | Propane Gas | Е | Trichloroethylene | Е |
| Isopropyl Acetate | Е | Propylene Glycol | Е | Triethylene Glycol | Е |
| Isopropyl Alcohol | Е | Propylene Oxide | Е | Tripropylene Glycol | Е |
| Isopropyl Benzene | Е | Resorcinol, Sat. | Е | Turpentine | Е |
| Kerosene | Е | Resorcinol, 5% | Е | Undecyl Alcohol | Е |
| Lactic Acid, 3% | Е | Salicylaldehyde | Е | Urea | Е |
| Lactic Acid, 85% | Е | Salicylic Acid, Powder | Е | Vinylidene Chloride | Е |
| Methoxyethyl Oleate | Е | Salicylic Acid, Sat. | Е | Xylene | Е |
| Methyl Alcohol | Е | Salt Solutions, Metallic | Е | Zinc Stearate | Е |

^{* =} Moderate Attack or appreciable absorption. Material will have limited life.

E = Excellent